

TOXIC CATCH:

Japan's unsustainable and
irresponsible whale, dolphin
and porpoise hunts



ABOUT EIA

EIA is an independent campaigning organisation committed to bringing about change that protects the natural world from environmental crime and abuse. As part of our work we have campaigned for three decades for effective protection for whales, dolphins and porpoises globally.

ACKNOWLEDGEMENTS

EIA would like to thank OceanCare and Wallace Global Fund for their support in making this work possible.

Report design by:
www.designsolutions.me.uk

October 2013

© Environmental Investigation Agency 2013

No part of this publication may be reproduced in any form or by any means without permission in writing from the Environmental Investigation Agency.



ENVIRONMENTAL INVESTIGATION AGENCY (EIA)
62/63 Upper Street
London N1 0NY, UK
Tel: +44 (0) 20 7354 7960
Fax: +44 (0) 20 7354 7961
email: ukinfo@eia-international.org
www.eia-international.org

EIA is solely and entirely responsible for the contents of this report.

COVER IMAGE:
© Erick Higuera

EXECUTIVE SUMMARY

Over a million toothed whales, dolphins and porpoises, commonly known as 'small cetaceans', have been killed in direct hunts in Japan in the past 70 years.

Catch limits set by the Government of Japan for 2013 permit the killing of 16,655 small cetaceans. This represents the largest directed hunt of cetaceans in the world. A comprehensive analysis of the available scientific data demonstrates unequivocally that there are grave concerns regarding the sustainability of these hunts.

Nine small cetacean species are targeted in Japan's coastal hunts, which take the form of small-type coastal whaling, hand harpoon hunts and drive hunts. Long before catch limits were introduced, the abundance of favoured species, such as the striped dolphin, began to drastically decline due to overexploitation.¹ As catches reached in excess of 30,000 small cetaceans per year concerns were raised at an international level regarding the unsustainable nature of Japan's hunts.² Catch limits were set by the Government of Japan in 1993; however, the actual catch numbers have declined to levels below the catch limits in the majority of species targeted by direct hunts. Declining demand for cetacean meat and the increasing economic costs of hunts may be playing a role, but there is significant evidence that a number of the exploited populations are depleted: changes in catch composition, declining abundance trends and reports from hunters of an increased difficulty in filling quotas all point to overexploitation.

Despite the indications of population declines, there appears to have been little monitoring of the status of the exploited small cetacean populations. For many of the species hunted, the last published abundance estimates are based upon surveys conducted more than 20 years ago.

Disregarding clear signals of overexploitation the Government has permitted catches to remain at levels that are unsustainable for eight of the nine target species.³ Small reductions in catch limits have been made since 2007 but often in prefectures where hunts are no longer occurring.

The Government of Japan provides little transparency regarding the methods it is using to set catch limits but they remain considerably higher than those that would be permitted under management strategies employed elsewhere in the world. In addition, there is little or no attention to catch composition or struck and lost rates - the latter remaining unaccounted for in the reported data on numbers killed. Although multiple tools now exist to calculate sustainable levels of marine mammal mortality these are not being employed.

The apparent absence of both up-to-date information on the status of populations and a scientifically rigorous method for setting catch limits displays a lack of responsibility by the Government to ensure the sustainability of small cetacean populations in Japanese waters. Through such conduct the Japanese Government is failing to implement its domestic policies of



sustainable utilisation⁴ and stipulations of the international conventions to which it is a signatory including the Convention on Biological Diversity.

In 2012 the Society of Marine Mammalogy, a professional society of more than 1,800 scientists from 60 countries expressed its concern regarding the sustainability of the hunts.⁵ In 2013, the Scientific Committee of the International Whaling Commission (IWC) again voiced its concerns regarding the sustainability of catches.⁶ However, the Japanese Government has continued to ignore IWC requests to reduce catch limits, persistently claiming that the IWC does not have competence with regards to small cetaceans.⁷

The hunts themselves serve only to provide toxic food products to Japanese consumers, who are largely unaware of the high levels of pollutants these top marine predators typically accumulate. Pollutant concentrations in meat and blubber from the marketplace can reach 85 times the safe limits for consumption of methyl mercury and 140 times the safe limit for PCBs.⁸ The Government of Japan's advisory limits remain wholly inadequate to inform or protect consumers – something all the more shocking in light of the recent signing of the Minamata Treaty in Japan.

Decades after concerns first arose, the Government of Japan continues to ignore international pleas to reduce catches, implement a scientific management system and publish up to date population assessments. Burying its head in the sand regarding the inevitable population declines and the health risks to Japanese consumers of whale, dolphin and porpoise products, the Japanese Government maintains a stubborn reluctance to relinquish this archaic industry for which there is declining domestic demand.

The Government of Japan has a responsibility to restore and maintain coastal cetacean species at their former levels, and protect consumers from the consumption of toxic seafood products. EIA urges the Government of Japan to phase out the hunts over a 10 year period, by establishing a scientific management programme that targets action on those species most at risk and working with hunters to find alternative livelihoods.



SMALL CETACEAN HUNTS IN JAPAN

© Mia Strickland/EIA



© David Sims/EIA

“Government catch limits have failed to adequately restrict hunts, and continue to permit hunts to operate at unsustainable levels.”

Three types of hunts target small cetaceans in Japan’s coastal waters: small-type coastal whaling, hand harpoon hunts and the drive hunts. Over the past century these hunts increased in scale and expanded their geographic range, capturing tens of thousands of small cetaceans every year. With the advent of faster motorised boats, hunts became increasingly efficient, resulting in the sequential depletion of small cetacean species in Japan’s national waters. In each type of hunt the same patterns of excessively high catches, overexploitation and subsequent declines in catches have been observed, concomitant with biological changes that signal the depletion of targeted populations. Government catch limits have failed to adequately restrict hunts, and continue to permit hunts to operate at unsustainable levels. As catches of favoured species such as the striped dolphin have fallen to unprecedentedly low levels, catches of less desirable species such as the bottlenose and Risso’s dolphin have increased and Japan’s Fisheries Agency has established quotas for new species.

SMALL-TYPE COASTAL WHALING

Small-type coastal whaling is conducted by small whaling vessels (less than 50 tonnes) which employ a canon below 50mm in calibre.⁹ These vessels target Baird’s beaked whales, short-finned pilot whales and false killer whales.¹⁰ Hunts occur for the most part within 50 nautical miles of the coast operating from whaling bases in Hokkaido, Miyagi, Chiba and Wakayama.¹¹

There has been a decline in overall small-type coastal whaling catches in the past decade, largely due to declining catches of both forms of the short-finned pilot whale, particularly those of the northern form. While the Government of Japan has failed to take action, whalers voluntarily deciding to ban hunts of the northern form since 2007.¹² Perhaps because of this, in 2012 the Government authorised an extension of the Baird’s beaked whale hunt through an expansion of permitted hunting seasons and an increase in the number of licensed vessels allowed to hunt in the Okhotsk Sea.¹³

DRIVE HUNTS

Drive hunts operate with a number of boats working together to locate and drive a group of dolphins towards an enclosed area. Noise created by banging 'trumpets' (long metal poles) on the vessels elicits a strong behavioural avoidance response, allowing fishermen to herd the dolphins over tens of kilometres to shore.¹⁴ Here they are entrapped with nets and may be left, thus confined, reportedly for as many as four days.¹⁵ Once the killing and live-capture begins, the nets are progressively tightened, confining animals into a small space in which they can be caught. Dolphins may be secured by their tail fluke and dragged by boats. Unable to control their surfacing to breathe during this period, dolphins sometimes die during the capture process due to forced drowning.¹⁶

Drive hunts were widespread along the coasts of the Sea of Japan, East China Sea and Pacific from the 14th until the late 19th/early-20th centuries.¹⁷ However by the mid-20th century drive hunts were limited to the Izu coast (Shizuoka prefecture), Taiji (Wakayama), Nago (Okinawa) and Nagasaki prefecture.¹⁸ The underlying causes of this geographical restriction in hunts remain unclear; declines in dolphin populations, declining demand for dolphin products and social changes in local communities are all potential factors.¹⁹

Although the number of villages hunting declined in the 20th century, the efficiency and potential range of hunts expanded, first in the 1920s with the advent of motorised vessels and again in the 1960s with improvements in vessel speed, allowing the capture of greater numbers of dolphins in a smaller amount of time.²⁰ Drive hunts were able to catch up to 21,000 dolphins annually (1942-1960), with captures predominantly comprised of striped dolphins.

Drive hunt captures peaked in 1959 when 21,953 striped dolphins were captured. Immediately following this catches of this species dropped significantly, by a degree of magnitude which cannot be explained by any decrease in hunting effort.²¹ Concomitant with catch declines that progressively extended across multiple prefectures, a decline in catch per unit effort of striped dolphins was observed, as well as a decrease in age of sexual maturity and an increase in the searching range required, all indicating population decline.²²

The introduction of limitations on the number of vessels and the months of drive hunt operation in the 1950s, and catch limits in the 1990s provided too little protection too late, with drive

hunts likely causing the complete eradication of the striped dolphin migration in Futo and significant declines elsewhere.²³ In response, when Wakayama catches have exceeded their catch limit the Japanese Government has permitted the transfer of the Chiba and Shizuoka quota for this species to Wakayama and Okinawa, allowing increased exploitation elsewhere in the species range. Hunts are now licensed to occur in two locations, Futo on the Izu coast in Shizuoka prefecture and Taiji in Wakayama prefecture, though only the latter currently conducts hunts.

The role of international markets

Drive hunts supply animals both for the live trade and as food products. While 99 per cent of the catches over the past 10 years have been for food, the increasing live dolphin trade is highly profitable and likely helps financially sustain the hunts for food in the face of declining demand for dolphin products. Over 1,500 small cetaceans have been captured in drive hunts and sold to aquaria globally over the past 26 years, with average numbers of live captures doubling from the 1990s to the 2000s. (Figure 1).²⁴ Live animals fetched between 660,000 and 7,712,000 yen (\$8,406 - \$98,222) per dolphin between 2002 and 2012. By comparison, a bottlenose dolphin killed for the meat trade will fetch 50,000 yen (approximately US\$500 as reported in 2009), less than a tenth of the value of a live dolphin.

International exports have exceeded \$15 million dollars over the past decade (2002-2012), with 389 dolphins transported to 15 different countries.²⁵ China has been the main importer, buying 248 dolphins (64 per cent of the exports). Sales also occur to aquaria within Japan but the number and value of these is not known. The global aquarium industry body, the International World Association of Zoos and Aquariums (WAZA), advises all WAZA member aquariums not to

BELOW:

Dolphins captured in Taiji drive hunts to be sold to aquaria or slaughtered for consumption.





© Pierre Gleizes/EIA

HAND HARPOON HUNTS

Hand harpoon hunts involve the pursuit of porpoises, dolphins and small whales at sea until they become exhausted and within reach for hunters to throw hand-held harpoons from the boats (EIA, 2000). Harpooned animals, often live, may then be left attached to flags or buoys while the hunters target additional animals.²⁶

Of the three types of hunts, harpoon hunts have the longest history, dating from prehistoric times.²⁷ In the 20th century they became large-scale operations, initially due to the introduction of fast motorised fishing vessels, followed by a second expansion in the 1980s, likely due to a declining whale meat supply following the cessation of Japan's commercial whaling and declining catches from the drive hunts due to over-hunting of striped dolphins.²⁸


Although eight different species can be taken in hand harpoon hunts, Dall's porpoise are the main target and have historically been hunted at unsustainable levels, with declining catches in recent years.²⁹ Over 400,000 Dall's porpoises have been killed in the hand-harpoon hunt since the moratorium on commercial whaling was implemented in 1986. Regulation of this hunt has been reported as inadequate, with problems in collation and calculation of catches leading to significant under-reporting of mortalities.³⁰

purchase dolphins which have been captured in the drive hunts.

Despite international criticism of the scale and killing methods of these hunts, the number of dolphins live captured and exported to international aquaria by Japan has increased. Kills are still an order of magnitude greater than live captures, but over the past decade the number of cetaceans killed has gradually declined while the number of live captures has increased (see Figure 1), in line with an increase in international exports. Hence, whilst the majority of drive hunts captures continue to be killed for meat, international markets for live captured animals appear to be playing an increasingly important role in the hunts.

FIGURE 1: Trends in live captures and kills in drive hunts in Taiji between 1986 and 2010

Source: National Research Institute of Far Seas Fisheries (NRIFSF, 1979-2011).





© Pierre Gleizes/EIA

CONSERVATION STATUS OF SMALL CETACEANS TARGETED IN COASTAL HUNTS

Cetaceans worldwide now face a litany of anthropogenic threats and those residing in Japan's coastal waters are no exception. Direct hunts alone have killed over a million small cetaceans in Japanese waters in the past 70 years. In addition to the significant levels of mortality from direct hunts, they suffer unknown levels of bycatch mortality in fisheries, habitat loss and degradation, prey depletion through overfishing, anthropogenic noise, vessel strikes, and increased pollutant loads. As such, the ability of populations to withstand the additional mortality caused by direct hunts is likely significantly reduced.

EIA's review of the status and threats faced by Japanese small cetaceans reveals worrying signs of overexploitation and depletion of a number of the local cetacean populations targeted by the hunts.³¹ Despite such signs, for the majority of species population assessments have not been regularly updated and catch limits remain set at unsustainable levels. For six of the nine species exploited the last population assessments were published more than 20 years ago. Although more recent surveys have been conducted for these species, these covered a much larger survey area and thus do not provide discrete population estimates of the populations exploited by hunts.³² Effort applied to coastal areas was low, limiting their accuracy and differences in season and survey area prohibit any analysis of population trends.³³

In 2013, EIA carried out a comprehensive analysis of the available data on the nine cetacean species targeted in direct

coastal hunts in Japan. The scientific paper was presented to the Scientific Committee of the International Whaling Commission (IWC) in June 2013, and is available at <http://iwc.int/sc65adocs>. The following is a summary of the main findings for each species.

DALL'S PORPOISE (*Phocoenoides dalli*)




© Luke Hyatt

There is a long history of unregulated exploitation of Dall's porpoises in Japan. This species has long formed the dominant component of the small cetacean catch in Japan, accounting for over 80 per cent of the catches from 1979-2010 (see Table 1).

There are two geographically segregated colour morphs of Dall's porpoises, the *dalli*-type and *truei*-type. Eight stocks have been identified, at least three of which are targeted by Japan's hunts.³⁴ These three stocks comprise one *truei*-type stock that breeds in the Central Okhotsk Sea, a *dalli*-type stock that breeds in the Southern Okhotsk Sea and

ABOVE:
Dall's porpoise laid out for auction at the fish market, Otsuchi.

FIGURE 2: Catches of Dall's porpoise



a third of unknown identity. The *truei* and *dalli*-type porpoises forming the majority of the catch are thought to originate from the Central and Southern Okhotsk sea stocks, respectively.³⁵

The most recent abundance estimates (see Table 2) are 23 per cent and 18 per cent lower than previous abundance estimates.³⁶ Abundance estimates are now ten years old and urgently require updating.

The Dall's porpoise hunt dates back to prehistoric time, with commercial hunting commencing in the 1920s and continuing to the present day. For much of this time hunts have been unregulated by catch limits, killing an average of more than 11,000 per year (1963-1992). In the 1980s the hunt expanded both geographically and seasonally. Kills peaked in 1988, with the hunt taking over 40,000 Dall's porpoises in a single year.³⁷ This has been attributed to a significant growth in demand for porpoise meat outside local hunting areas, generated by a reduction in availability of minke whale meat once the moratorium on commercial whaling was implemented. Catches have since declined steeply (see Table 1) and ceased entirely during 2011 due to the Great East Japan Earthquake and tsunami (see Figure 2).³⁸ The hunt resumed in March 2012, and approximately 1,200 individuals were landed in Iwate during the November 2012 to April 2013 season.³⁹

While historic catch levels have been alarmingly high and significantly exceeded sustainable levels, published data have also historically significantly under-reported actual mortalities.⁴⁰ Incomplete collation of records and inaccurate calculation of catches, is estimated to have led to under-reporting of mortalities by up to 11,000 animals

per year, while failure to take into account struck and lost takes (where the animal is struck by the harpoon but not landed), has been estimated to result in total mortality 10-14 per cent higher than the number landed.⁴¹

Trends in catch composition have indicated a worrying increase in the proportion of mature and lactating females being caught in the Sea of Japan.⁴² This removal of mature lactating females further reduces the recruitment potential of the population, and at the same time likely results in the mortality of dependent calves. Given the evidence of female philopatry it poses the additional risk of localised depletion. Although operational months are limited to certain months by prefecture, the Dall's porpoise hunts operate at some locations throughout the year (starting on August 1 and closing on July 31) and target the species in sensitive breeding and calving periods.⁴³

Catch limits introduced in 1993 were based on an annual Allowable Biological Catch (ABC) of approximately 4 per cent of the abundance estimate.⁴⁴ This method of setting catch limits permits catches much higher than thresholds used to manage cetacean populations in other regions of the world and fails to take into account potentially significant levels of mortality from bycatch or other causes. For example, catch limits for the 2013/14 season are still 4.8 and 4.7 times higher for the *dalli* and *truei*-type than the 'robust' Potential Biological Removal (PBR) threshold, a threshold designed to prevent populations from declining below their maximum net productivity level.⁴⁵ The catch limits equate to 4.1 per cent (*dalli*-type) and 3.9 per cent (*truei*-type) of the 2007 abundance estimates. If assumptions on population growth rates are correct, allowing a catch of more than 4 per cent will cause the population to decline to levels approaching zero, and will prevent future recovery.⁴⁶

Japanese Government scientists have fully recognised the inadequacy of their management approach, stating that "... the current management procedure, based on only the best values, could fail to manage the stocks at a considerably high probability".⁴⁷ Although there have been repeated calls from the IWC Scientific Committee to reduce catch limits to sustainable levels more than 20 years on this has not occurred.⁴⁸

STATUS: Unknown, some populations likely depleted.

BAIRD'S BEAKED WHALE (*Berardius bairdii*)



© Adam James Searcy

Three populations of Baird's beaked whales are hunted off the coast of Japan, one off the Pacific coast, one in the eastern Sea of Japan and a third in the southern Okhotsk Sea. Abundance estimates for two of the three populations are more than 20 years old – their status is therefore unknown.

Hunting of Baird's beaked whales began around 1600, primarily in the seas around Chiba prefecture in hand harpoon hunts.⁴⁹ Hunting by small-type coastal whaling vessels began in the early 20th century and increased sharply following World War II, spreading to Hokkaido, the north-east (Sanriku) coast and to the Sea of Japan.⁵⁰ Hunts now occur primarily on the Pacific coast and in the Okhotsk Sea.⁵¹

The IWC small cetacean sub-committee has noted that the current catch level "represents about one per cent of the estimated population size and in the absence of an estimate of gross reproductive rate, was unable to determine whether or not the population could sustain the catches".⁵² Reiterating previous advice⁵³, in 2012 the Scientific Committee therefore recommended that: "(1) It is especially important to clarify population structure and geographical boundaries of the stocks off Japan, particularly as long as hunting continues there; (2) Improved and updated abundance estimates are needed for each population, and trends in abundance should be assessed. These needs particularly apply to exploited stocks".⁵⁴

The annual catch limit has been increased several times since first introduced and currently totals 66 whales (see Figure 3). In 2012, the hunting season and number of vessels licensed to hunt Baird's beaked whales in specific areas was increased.⁵⁵ The reason for this increase in effort is unclear as allocated quotas are already largely being filled.

STATUS: Unknown.

FIGURE 3: Catches of Baird's beaked whales





FIGURE 4: Catches of short-finned pilot whale



SHORT-FINNED PILOT WHALE (*Globicephala macrorhynchus*)



© WDC

Two forms of the short-finned pilot whale occur off Japan, known as northern and southern forms. The taxonomic status of these two forms is unresolved. The two forms may represent separate species or subspecies but are currently treated as a single taxonomic unit.⁵⁶ If they are separate species, then each may warrant being listed as threatened according to IUCN criteria.⁵⁷

FIGURE 5: Catches of Risso's dolphin





FIGURE 6: Catches of striped dolphin



As far back as the 1980s changes in catch composition of the northern form indicated overhunting and population decline. This has prompted the IWC Scientific Committee to state that “it was desirable that no animals be taken until we have a clearer understanding of the status of the stock”.⁵⁸ Catches continued regardless and recent analyses now indicate a drastic population decline in the 1980s when catches far exceeded the PBR threshold. There has been a slight recovery since but not to former levels and Japanese scientists conclude that whaling may have “seriously depleted the abundance of the north form”.⁵⁹

The annual catch limit continues to be set above the PBR threshold providing little protection from continued overexploitation. Compensating for the inaction of the Government of Japan, whalers have implemented a voluntary ban on hunting of the northern form since 2007.⁶⁰

Catches of the southern form have also declined, reaching their lowest in 2010 (see Table 1 and Figure 4). Current catch limits remain above sustainable levels, risking further population depletion (see Table 2).⁶¹

STATUS: Northern form significantly depleted, status of the southern form unknown; both forms assessed as rare by the Mammalogical Society of Japan.

RISSO'S DOLPHIN (*Grampus griseus*)



© Jim Scarff

The Japanese drive hunts, small-type coastal whaling and hand harpoon hunts have all regularly hunted Risso's dolphins, both for food and for live captures to supply the aquarium industry.

Catches increased during the 1990s, perhaps in response to the decline in more popular species, with approximately 200-500 Risso's dolphins killed per year (1993-2010) (see Table 1 and Figure 5). Although there have been recent reductions in catch limits they are still almost double (1.9 x PBR threshold) sustainable levels (see Table 2),⁶² and remain inadequate to protect the population(s) from depletion. Published abundance estimates of hunted populations are now over 20 years old and urgently require updating. The status of exploited populations is therefore unknown.

STATUS: Unknown.

STRIPED DOLPHIN (*Stenella coeruleoalba*)



© M Pilot

For over 100 years striped dolphins have been heavily hunted in Japanese waters in drive and hand-harpoon hunts.⁶³ Prior to the introduction of quotas, over

159,500 striped dolphins were killed between 1963 and 1992 and it is thought that by the 1990s the Japanese drive hunt “had depleted coastal stocks of striped dolphins to less than 10 per cent of the post-World War II level”.⁶⁴

On the Izu coast (Futo and Kawana) hunts may have led to the complete eradication of the local striped dolphin population.⁶⁵ Declines extend across multiple prefectures; in addition to the lack of catches in Futo and Kawana, catches in Taiji fell by more than 90 per cent (1980-1991) despite an increase in the number of searching vessels over the same period.⁶⁶ Similarly in Chiba, striped dolphins have not been successfully hunted since 1995.


In addition to the long-running decline in catches of striped dolphins, reproductive parameters have changed in a manner consistent with a density decline. A decreased age of sexual maturity, characteristic of a density-dependent response to population reduction has been observed, as well as a decline in catch per unit effort and an increase in the searching range required, all indicating population decline and overexploitation of this species.⁶⁷

In response to this overwhelming evidence, the Japanese Government has only made reductions in the catch limit in those prefectures where catches are no longer taking place, imposing no reduction on actual catches. In fact, when Wakayama hunters have exceeded their catch limits the Japanese Government has permitted the temporary transfer of the Chiba and Shizuoka quota to Wakayama and Okinawa, allowing an increase in exploitation of the species elsewhere in its range.

Current catch limits remain 5.3 times higher than the sustainable level based on a PBR threshold, and average actual catches (2007-2011) are 4.4 times the PBR threshold, far exceeding sustainable levels (see Table 2).⁶⁸ The published abundance estimates upon which we presume such catch limits are based are now over 20 years old, and with current abundance and population structure remaining unresolved, updated assessments are urgently required.

STATUS: Endangered, some populations may be locally extinct.

FIGURE 7: Catches of spotted dolphin



PAN-TROPICAL SPOTTED DOLPHIN (*Stenella attenuata*)



© Mike Grimes, Oretani Wildlife, www.oretani.com

Spotted dolphins are killed in large numbers by drive hunts in Japan. Declines in catches of the pan-tropical spotted dolphin occurred in the late 1980s and early 1990s (see Figure 7). Recently annual catches have continued to decline, alongside which there has been a possible decline in the minimum age of sexual maturity in females and a decline in catch per unit effort, indicative of an abundance decline in the local population(s).⁶⁹

Annual catch limits were reduced in 2007, but only in those prefectures where catches are no longer taking place, imposing no reduction on current catches. Despite the signals of overexploitation, catch limits remain considerably higher than any catch attained since 1988 and are far above sustainable levels, being 5.7 times higher than a PBR threshold (see Table 2).⁷⁰

Published estimates are now 20 years old and stock structure remains unresolved – further studies are urgently required to assess hunted populations’ status.

STATUS: Likely declined in abundance, status unknown.

FIGURE 8: Catches of bottlenose dolphin





FIGURE 9: Catches of false killer whale



COMMON BOTTLENOSE DOLPHIN (*Tursiops truncatus*)



© Sarah Baulich

The bottlenose dolphin is the main species targeted in the drive hunts for the live capture trade, comprising 76 per cent of the live captured individuals since records of live captures began.⁷¹ They are also killed for consumption and have previously been culled in Japanese waters due to claims of competition with fisheries. In the late 1970s-1995 several hundred were culled annually off Iki Island and the Kii Peninsula.⁷²

In the 1980s the numbers killed in hunts increased to more than 900 per year, likely far exceeding sustainable levels. Catches have since declined but catch limits remain more than double a PBR threshold (see Table 2 and Figure 8).⁷³

Meanwhile the Government has failed to publish up to date abundance estimates for the exploited population(s). As with other species, these are now more than 20 years old, and further surveys are therefore urgently required.

STATUS: Unknown, assessed as threatened by the Mammalogical Society of Japan.

FALSE KILLER WHALE (*Pseudorca crassidens*)



© Bo Pardaou

Historically the false killer whale appears to have been relatively common off the Japanese coast but with abundance estimates of the population(s) targeted by hunts now more than twenty years old, knowledge of their status urgently requires updating.

Catches were highest in the 1970s and 1980s, peaking at 356 in 1980. As well as being hunted for consumption, they are live-captured and have previously been killed for their depredation of fisheries. Over 900 were culled between 1965 and 1980 around Iki Island in Japan due to interactions with the yellowtail fishery.⁷⁴

Following the high number of kills in the 1970s and 1980s catches declined and from 1986 to 1992 did not exceed 100 per year with many years of zero, or near-zero, catches (see Figure 9). Although targeted by all three hunts they are currently captured in relatively small numbers, most likely due to low encounter rates.

Despite catches not even approaching the previous catch limit of 50 per year, the catch limit was increased in 2007 and there is now a total catch limit of 120 for this species (see Figure 9). Such catch limits are clearly far above likely sustainable levels, being 8.6 times the PBR threshold and equating to 5.9 per cent of the abundance estimate (see Table 2). Such levels are particularly irresponsible given that it is a species which typically occurs with low abundance, about which relatively little is known and which may be vulnerable to even low-level threats.⁷⁵

STATUS: Unknown, assessed as threatened by the Mammalogical Society of Japan.

PACIFIC WHITE-SIDED DOLPHIN (*Lagenorhynchus obliquidens*)




© Luke Hyatt

High catches of several thousand Pacific white-sided dolphin per year occurred in 1983 and 1984. Subsequently catches declined and remained below 50 per year.

Hunts for consumption ceased in 1993 as a quota was not created for this species but live captures continued. In 2007 a new quota of 360 dolphins was established for the species (see Figure 10).⁷⁶ This new quota could represent an effort to compensate for declining catches of other species or may also be due to demand for this species by the aquarium industry.⁷⁷

FIGURE 10: Catches of Pacific white-sided dolphin



The current catch limit is slightly higher than the PBR threshold (see Table 2). With abundance estimates now more than 15 years old and with evidence of significant sub-population structure, the status of the population(s) being targeted by Japanese hunts is unknown.


STATUS: Unknown.

TABLE 1. Reported catches by species, 1986-2011⁷⁸

Year	Baird's beaked whale	Bottlenose dolphin	Dall's porpoise	False killer whale	Pacific white-sided dolphin	Risso's dolphin	Short-finned pilot whale	Spotted dolphin	Striped dolphin	Total
Total catches reported before 1986	5,290	8,191	190,167	1,303	5,208	1,678	8,634	21,265	323,804	N/A
1986	40	238	16,515	4	0	0	376	891	2,770	20,834
1987	40	1,810	25,600	33	0	6	386	1,815	389	30,079
1988	57	812	40,367	70	0	124	571	1,879	2,227	46,107
1989	54	390	29,048	30	0	14	252	129	1,225	31,142
1990	54	1,298	21,804	81	39	93	167	11	749	24,296
1991	54	433	17,634	54	12	394	355	153	1,022	20,111
1992	54	173	11,403	97	0	121	360	636	1,045	13,889
1993	54	215	14,318	20	1	505	337	565	544	16,559
1994	54	362	15,947	0	6	312	196	449	545	17,871
1995	54	963	12,396	49	2	405	239	105	539	14,752
1996	54	314	16,100	40	19	372	484	67	303	17,753
1997	54	352	18,540	43	0	228	347	23	602	20,189
1998	54	266	11,385	48	0	445	229	460	449	13,336
1999	62	749	14,807	5	11	489	396	38	596	17,153
2000	62	1,426	16,171	8	1	506	304	39	300	18,817
2001	62	259	16,650	37	6	477	391	10	484	18,376
2002	62	801	15,949	7	7	387	176	418	642	18,449
2003	62	180	15,720	21	8	379	160	132	450	17,112
2004	62	652	13,789	7	12	511	176	2	637	15,848
2005	66	361	14,664	1	0	394	176	13	457	16,132
2006	63	372	12,014	35	0	344	271	405	515	14,019
2007	67	401	11,357	4	0	517	338	16	470	13,170
2008	64	391	7,226	5	21	338	181	329	598	9,153
2009	67	433	9,540	1	21	430	295	3	419	11,209
2010	66	434	4,919	0	27	397	44	132	558	6,577
2011	61	119	1,952	20	24	377	126	108	502	3,289
TOTAL	6,793	22,395	595,982	2,023	5,425	10,243	15,967	30,093	342,841	1,031,762

TABLE 2. Recent quotas and catches in relation to abundance data and sustainable thresholds.⁷⁹

Species	Estimated abundance CV	Survey period	Area surveyed	Catch limit (2013-2014)	Average annual catch (2007-2011)	PBR threshold (0.5) (Funahashi & Baker, 2011)	Catch limit as a % of abundance
Dall's porpoise (<i>truei</i>)	217,000 CV=0.227 ⁸⁰	1989-1990	40°N-51°N, 140°E-170°E	6,656	1,956	-	-
	178,157 CV=0.23 ⁸¹	2003	Central Okhotsk Sea			1,472 ⁸²	3.7%
Dall's porpoise (<i>dalli</i>)	226,000 CV=0.154 ⁸³	1989-1990	40°N-51°N, 140°E-170°E	6,837	5,042	-	-
	173,638 CV=0.21 ⁸⁴	2003	Southern Okhotsk Sea			1,483 ⁸⁵	3.9%
Baird's beaked whale	4,200 CV=0.295 ⁸⁶	1984	Pacific coast	52	65 (for Pacific, Sea of Japan & Okhotsk Sea combined) ⁸³	-	1% ⁸⁴
	3,950 CV=0.28 ⁸⁷	1983-1989					
	5,029 95% CI = 1,801-14,085 ⁸⁸	1991-1992					
	10,190 ⁸⁹	2008					
	7,307 ⁹⁰	2009					
	1,260 CV=0.45 ⁹¹	1983-1989	Sea of Japan	10		-	0.8%
	660 CV=0.27 ⁹²	1983-1989	Okhotsk Sea	4		-	0.6%
Short-finned pilot whale (northern form)	5,344 ⁹⁵	1984-1985	Unknown	36	0	-	-
	4,239 CV=0.61 ⁹⁶	1986-1988	Unknown			-	-
	3,879 CV=0.49 ⁹⁷	2006	Southern and northern coast of Sanriku & southwest coast of Hokkaido			26 ⁹⁸	0.9%
Short-finned pilot whale (southern form)	14,012 CV=0.229 ⁹⁹	1983-1991	north of 30°N, west of 145°E	219	197	116	1.6%
Risso's dolphin	31,012 CV=0.211 ¹⁰⁰	1983-1991	north of 30°N, west of 145°E	487	412	260	1.6%
Striped dolphin	19,631 CV=0.696 ¹⁰¹	1983-1991	north of 30°N, west of 145°E	595	509	116	3%
Spotted dolphin	15,900 CV=0.401 ¹⁰²	1983-1991	north of 30°N, west of 145°E	606	118	115	3.8%
Bottlenose dolphin	36,791 CV=0.250 ¹⁰³	1983-1991	north of 30°N, west of 145°E	673	356	299	1.8%
False killer whale	2,029 CV=0.429 ¹⁰⁴	1983-1991	north of 30°N, west of 145°E	120	6	14	5.9%
Pacific white-sided dolphin	56,764 CV=0.8 ¹⁰⁵	1992-1996	north of 30°N, west of 145°E	360	19	314	0.6%

A large whale, likely a humpback whale, is shown swimming underwater in clear blue water. The whale's head and upper body are visible, with sunlight filtering through the water, creating shimmering patterns on its dark skin. The whale is moving towards the left of the frame. The water is a deep, vibrant blue, and the surface of the water is visible at the top of the image.

“ Catch limits for all species bar one are set above sustainable levels. Published abundance estimates for many populations have not been updated for over 20 years.”



© Pierre Gleizes/EIA

THE INDIRECT IMPACTS OF HUNTING

The officially reported catches of cetaceans in Japanese waters are alarmingly high and are widely considered unsustainable.¹⁰⁶ But added to these reported catches are additional mortalities, never monitored or reported, with likely population-level effects that go beyond individual mortalities. The unknown impacts of stress and social disruption brought about by hunts are likely to reduce survivorship and reproductive success of the remnant populations and impede their recovery. This exacerbates a low baseline rate of recovery that results from odontocetes' life history, social and behavioural characteristics.¹⁰⁷ Some of the populations targeted by Japan's hunts are already showing signs of significant declines in abundance to a point where they may no longer have the capacity to recover.

UNDOCUMENTED DEATHS

In the past, published figures have significantly under-reported the landed catches of Japan's small cetacean hunts. Examination has focused on the Dall's porpoise hunt but the incomplete reporting may also extend to catch statistics for other small cetacean species taken by hand harpoon hunts.

In addition to the widespread under-reporting of landed catches, other mortalities are never reported in official figures for any of the three types of hunts. These include struck and lost individuals, deaths of dependent juveniles, and until 1986, live captures, although clearly these all remove individuals from wild populations.

In the Dall's porpoise hand harpoon hunt, struck and lost individuals were previously estimated to result in a total mortality 10-14 per cent higher than the number landed,¹⁰⁸ whilst in the drive hunt there are known juvenile deaths and those that die 'naturally' in the enclosure, which are discarded and not considered part of the catch.¹⁰⁹

POPULATION LEVEL IMPACTS OF STRESS?

Individually and cumulatively, Japan's drive, hand harpoon and small-type whaling hunts repeatedly disturb targeted populations. Vessel noise and the pursuit itself are likely to induce stress not only in animals selected for killing but, more significantly for the long-term conservation of populations, also in non-target individuals in the immediate social group, other groups in the wider vicinity of hunts and animals captured and then released. Hunts may therefore induce stress in and disturb and displace non-target individuals over a wide geographic area.¹¹⁰

The drive hunts in particular regularly pursue dolphins for extended periods of time but ultimately fail to catch them. The pursuit phase may cause stress-induced pathology that can lead to disease and unobserved mortality in animals that are not killed in the hunt.¹¹¹ Some animals are also released after being confined and subjected to

acute noise and stress. Lethal and sub-lethal stress-induced changes in released individuals add to the known high levels of mortality and may reduce the reproductive potential of remnant populations and their ability to recover from such removals, with potential population level consequences. Due to their relatively old age of first reproduction and low calving rate, odontocete populations can be overexploited by catches of only a few percent of the total population per year, and are less resilient to overexploitation than other species groups.¹¹² Adding to this, the highly social structure of odontocete societies means that social disruption caused by exploitation may reduce survivorship and reproductive potential of the remaining population, further impeding their ability to recover. Studies now indicate a lack of strong recovery in other heavily exploited odontocete populations even decades after intense exploitation has ceased.¹¹³

There are therefore multiple factors likely to be impairing the recovery Japan's small cetacean populations, even in the absence of continued exploitation. An up-to-date assessment of the status of exploited species and the development of sustainable catch limits which take into account the multiple anthropogenic and environmental threats these populations are facing and their inherent capacity to recover are urgently required in order to prevent further declines and the potential for localised extinctions.

BELOW:
Pacific white-sided dolphin
in the wild.



© Luke Hyatt



© Paul Redman/EIA

HUMAN HEALTH RISKS

“Levels of pollutants in tissues of small cetaceans from Japanese waters have far exceeded the advisory limits for human consumption, with concentrations of total mercury that are more than 200 times Japan’s limit.”

During the past 100-200 years, concentrations of a number of chemical pollutants have increased dramatically in the marine environment. The toxicological effects of such pollutants pose a global threat to the health and viability of cetacean populations and the health of human populations consuming them. Mercury and persistent organic pollutants (POPs) bioaccumulate and biomagnify in the food chain.¹¹⁴ As many cetacean species feed at a high trophic level and are long-lived they can accumulate high doses of POPs and heavy metal pollutants to the extent that concentrations of POPs can reach levels 70,000 times higher than background environmental levels.¹¹⁵

The high pollutant levels that have been documented in Japanese small cetaceans are a further threat to cetacean populations already beleaguered by hunting, but also a significant health threat to the human populations that consume their meat and blubber. In cetaceans, POPs have been linked to increased rates of cancer, increased first calf mortality, immune suppression and a higher susceptibility to infectious disease.¹¹⁶ At a population

level, they have been postulated to be a primary factor causing population declines and suppressing growth and recovery of populations.¹¹⁷

The consumption of cetacean products contaminated with high levels of POPs and mercury poses a grave health risk to humans. Ingestion of these toxins have been linked to a range of immunological, cardiovascular and reproductive effects in humans, including impaired foetal neurological development, and an increased risk of Parkinson’s disease, arteriosclerosis and diabetes.¹¹⁸

Levels of pollutants in cetacean meat being sold for consumption have far exceeded the advisory limits for human consumption, with concentrations of total mercury that are more than 200 times Japan’s limit of 0.4 parts per million of mercury (ppm); methylmercury that is 26 times higher than the WHO limit of 1.0ppm for large predatory fish and 87 times higher than Japan’s limit of 0.3ppm (although this excludes predatory fish); and PCB levels that are more than 75 times higher than Japan’s limit of 0.5ppm.¹¹⁹ Based on samples of

boiled whale internal organs that displayed such mercury levels, studies in rats have indicated that, irrespective of chronic effects, a single ingestion of boiled whale internal organs may cause acute intoxication by inorganic mercury.¹²⁰

Such high levels of pollutants do not appear anomalous. What is deeply concerning is that across large numbers of samples, average concentrations of mercury exceed advisory limits in all eight species tested.¹²¹

Concern has been raised since the release of radioactive material from Fukushima but there appears to be no monitoring of levels in small cetacean species, although they can bioaccumulate high levels of radioactive elements.¹²² The Dall's porpoise is one species whose distribution may overlap with the fallout of radioactive caesium from Fukushima. Levels in prey species of the Dall's porpoise have exceeded the Government of Japan's safety limit of 100Bq/kg in 2010-2012 and levels in cetaceans could be an order of magnitude higher - studies in the Barents and Norwegian Sea have shown that concentrations of 137 Cs were 10 fold higher in the harbour porpoise (*Phocoena phocoena*) than in species at the lower levels of the food web.¹²³ As such it is possible that cetaceans are bioaccumulating very high levels of radioactive elements, presenting a severe risk to consumers, as well as a novel threat to cetaceans. Fish, molluscs and crustaceans offshore from Fukushima have received doses that are likely to have reproductive effects and may markedly increase mortality.¹²⁴ The levels and impact in small cetaceans remain unknown, with no testing of small cetacean species according to the Government of Japan's online database.¹²⁵

AN INADEQUATE RESPONSE - JAPAN'S GUIDELINES AND MONITORING FOR POLLUTANTS

In reaction to the Minamata disease tragedy, the Japanese Ministry of Health and Welfare set provisional permitted levels of T-Hg and M-Hg in marine foods at 0.4 and 0.3ppm respectively, however these limits are not applied to cetacean products or indeed other predatory fish species that are likely to exhibit high levels.¹²⁶ As such, to EIA's knowledge, the sale of highly contaminated meat continues unmonitored and unregulated.

Advice on consumption limits is limited to pregnant women and covers only a subset of the cetacean species consumed by Japanese citizens, excluding several species that are known to be highly contaminated. Japan's guidance remains far more limited and recommends far higher levels of 'safe' consumption than that provided to populations that consume similarly polluted cetacean products in other parts of the world. Indeed, in the Faroe Islands medical authorities have recommended that due to the health implications, pilot whale should no longer be consumed at all.¹²⁷

Monitoring for health effects in Japan is as inadequate as the advice given to consumers. Evidence of adverse effects from the consumption of polluted cetacean meat is now accumulating from other countries where aboriginal subsistence whaling occurs and includes multiple immunological, cardiovascular and reproductive effects.¹²⁸ In 2010, the National Institute of Minamata Disease carried out a study to measure mercury levels in the hair of Taiji residents. The tests showed that average mercury levels in Taiji were higher than the national average, with a number of individuals with levels above the World Health Organisation (WHO) limit for neurological effects (50ppm), something that has not been observed elsewhere in Japan in recent tests. However no action has since been taken to reduce consumption of small cetaceans in Taiji.¹²⁹

BELOW:

This packet of whale giblets purchased from Amazon.co.jp was tested in a Japan-based laboratory and found to contain 21ppm mercury, over 50 times higher than the regulatory limit. Amazon.co.jp subsequently banned the sale of all whale, dolphin and porpoise products. .





CONCLUSIONS AND RECOMMENDATIONS

Over a million small cetaceans have been killed in direct hunts in Japan in the past 70 years. Over the decades, a number of problems have been identified with the management of small cetacean hunts in Japan, which remain largely unresolved. This report shows that:

- There is a lack of transparency regarding methods used to set catch limits;
- Catch limits for all species examined (i.e. excluding Baird's beaked whale) are set considerably above likely sustainable levels, based on calculations using a PBR threshold (see Table 2), even where this is calculated assuming only moderate exploitation and no other sources of anthropogenic mortality;¹³⁰
- With the exception of some populations of Baird's beaked whales and the northern-form short finned pilot whale, abundance estimates have not been regularly updated, with most more than 20 years old whereas best practice requires surveys every eight years;¹³¹
- There is a lack of regulation and enforcement of catch limits including extended months of operation when quotas could not be filled and transfer of quotas between prefectures when catches have exceeded local limits;
- There has been incomplete collation of catch data, and no regular reporting or estimates of struck and lost rates or other known mortalities.

The Government of Japan claims to support a policy of sustainable utilisation of marine resources and stipulates that fisheries should be based on sustainable principles.¹³² As a signatory to the international Convention on Biodiversity (1992), Japan is committed to take measures to ensure the conservation and sustainable use of biodiversity. The Government of Japan justifies supporting cetacean hunts on the grounds of "sustainable use of marine living resources" and "the principle of science-based management of resources".¹³³ Despite such claims and commitments, the Government of Japan continues to set catch limits at levels that are highly unsustainable and has failed to carry out adequate monitoring of populations which are being exploited.

In expert international fora such as the IWC Scientific Committee and the Society of Marine Mammalogy, concerns have been raised on numerous occasions over the status of populations targeted by Japan's hunts.¹³⁴ Japan has failed to respond, claiming only that the IWC does not have competence with regards to small cetaceans.

The apparent continued use of an outdated method for calculating catch limits, or for some species perhaps no scientific method at all, as well as the lack of up-to-date published abundance assessments undermines the sustainable management stipulated in Japan's domestic laws and the international laws to which it is a signatory. With the exception of only one species, quotas continue to be set above sustainable levels.¹³⁵ Even the warning signs of overexploitation and population decline that have been observed in four of the nine species hunted have not

stimulated adequate adjustment of quotas or prompted reassessment of populations' status. There are multiple causes for concern regarding the likelihood of recovery of Japan's small cetacean populations, even in the absence of continued exploitation. Management must become more precautionary if further declines are to be prevented.

An up-to-date assessment of the status of exploited species and the development of a scientific management approach which takes into account the multiple anthropogenic and environmental threats these populations are facing and their inherent capacity to recover is urgently required in order to prevent further declines and the potential of localised extinctions.

EIA urges the Government of Japan to:

- Respond to IWC requests and immediately suspend hunts of species showing the most severe signs of over-exploitation, in particular the Dall's porpoise, northern form of the short-finned pilot whale and striped dolphin;
- Conduct an up-to-date assessment of the status of all species taken by hunts, including studies of population structure;
- Collect and publish data on struck and lost rates, bycatch, hunt effort and reproductive status, sex and age composition of catches;
- Monitor targeted small cetacean populations for stress-induced impacts and the effects of social disruption;
- Reform the management strategy to bring it in line with modern international conservation management strategies, taking into account other mortalities such as struck-and-lost individuals, bycatch and other environmental/anthropogenic threats. This should be based upon up-to-date knowledge of population status and intrinsic recovery rates;
- Introduce independent observation of landed catches of all hunts and enforce any breaches with penalties;
- Establish and enforce time and area restrictions on hunts in order to protect cetacean species during sensitive breeding and calving periods;
- Conduct and publish long-term monitoring studies of pollutant levels in cetaceans;
- Phase out all small cetacean hunts over a ten year period, starting with those populations most at risk and those demonstrating the highest levels of pollutants.

REFERENCES

1. Kasuya, T. 1999a. Review of the biology and exploitation of striped dolphins in Japan. *Journal of Cetacean Research and Management* 1(1): 81-100.
2. IWC, 1992a. Annual Report of the International Whaling Commission 42:182-221.
3. Funahashi, N. & Baker, C. S. 2011. Japan's hunting of dolphins - Is it sustainable? Presentation at the 2011 Society for Marine Mammalogy conference, Kanaji, Y., Okamura, H. and Miyashita, T. 2011. Long-term abundance trends of the northern form of the short-finned pilot whale (*Globicephala macrorhynchus*) along the Pacific coast of Japan. *Marine Mammal Science* 27:477-492.
4. Wade, P. R., Bass, C. L. & Kasuya, T. 2008. A comparison of methods for calculating thresholds for evaluating levels of catch of Japan's Dall's porpoise (*Phocoenoides dalli*) hand-harpoon hunt. SC/60/SM24 presented to 60th annual meeting of the IWC Scientific Committee in Santiago, Chile.
5. Fisheries Basic Act, 2001 Available at <http://faolex.fao.org/BasicAct/Biodiversity>, 2008 Available at <http://faolex.fao.org/SMM>, 2012 Letter to Japanese Government. Available at: http://www.marinemammalscience.org/index.php?option=com_content&view=article&id=675:additional-letter-to-japanese-government-regarding-dolphin-and-small-whale-hunts&catid=65:presidential-letters&Itemid=183
6. IWC, 2013. Report of the Scientific Committee, unpubl. Available online at: <http://iwc.int/scientific-committee-reports>.
7. IWC, 2002a. Annual Report of the International Whaling Commission 54: 40.
8. Endo, T., Haraguchi, K., Hotta, Y., Hisamichi, Y., Lavery, S., Dalebout, M. and Baker, C. S. 2005. Total mercury, methyl mercury and selenium levels in the red meat of small cetaceans sold for human consumption in Japan. *Environmental Science and Technology* 39, 5703-5708.
9. Kasuya, T. 2007. Japanese Whaling and other cetacean fisheries. *Environmental Science and Pollution Research* 10: 39-48.
10. Fisheries Research Agency, 2012a. The State of International Fishing Resources in 2011 - Small Cetaceans
11. Ref 9, *ibid*.
12. Kanaji, Y., Okamura, H. and Miyashita, T. 2011. Long-term abundance trends of the northern form of the short-finned pilot whale (*Globicephala macrorhynchus*) along the Pacific coast of Japan. *Marine Mammal Science* 27:477-492
13. Fisheries Research Agency, 2012b. The State of International Fishing Resources in 2011 - Baird's beaked whales.
14. Akamatsu, T., Hatakeyama, Y. and Takatsu, N. 1993. Effects of pulse sounds on escape behaviour of false killer whales. *Bulletin of the Japanese Society of Scientific Fisheries* 59(8): 1297-303. Brownell, R., Nowacek, D. & Ralls, K. 2008. Hunting cetaceans with sound: A worldwide review. Publications, Agencies and Staff of the U.S. Department of Commerce. Paper 94.
15. Anon, pers. comm.
16. Butterworth, A., Brakes, P., Vail, C. S. & Reiss, D. 2013. Veterinary and Behavioral Analysis of Dolphin Killing Methods Currently Used in the "Drive Hunt" in Taiji, Japan. *Journal of Applied Animal Welfare Science* 16(2).
17. Ref 9, *ibid*.
18. Ref 9, *ibid*.
19. Ref 9, *ibid*.
20. Ref 9, *ibid*.
21. Ref 9, *ibid*.
22. IWC, 1993a. Report of the scientific committee 43: 82-85.
23. Kasuya, T. 2007. Japanese Whaling and other cetacean fisheries. *Environmental Science and Pollution Research* 10: 39-48.
24. National Research Institute of Far Seas Fisheries (NRIFSF, 1979-2011). Japanese Progress Reports on small cetacean research. Reports from 1979-1998 are published within IWC annual reports, SC/32-SC/52. Those from 1999-2010 are available online at: http://www.jfa.maff.go.jp/j/whale/w_document/
25. Ref 9, *ibid*.
26. National Research Institute of Far Seas Fisheries (NRIFSF, 1979-2011). Japanese Progress Reports on small cetacean r research. Reports from 1979-1998 are published within IWC annual reports, SC/32-SC/52. Those from 1999-2010 are available online at: http://www.jfa.maff.go.jp/j/whale/w_document/
27. Ref 9, *ibid*.
28. Ref 9, *ibid*.
29. Ref 24, *ibid*. IWC, 2001a. Annual Report of the International Whaling Commission, 53: 30-32.
30. Fujise, Y., Ishikawa, H., Saino, S. & Kawasaki, M. 1993. Catch and struck-and-lost rate in the Japanese Dall's porpoise hand-harpoon fishery. SC/44/SM18 in Report of the International Whaling Commission 43: 453-457.
31. Kasuya, T. 1992. Examination of Japanese statistics for the Dall's porpoise hand harpoon fishery. Report of the International Whaling Commission 42, 521-528.
32. Baulch, S. & Perry, C. 2013. Status review of small cetacean species targeted by direct hunts in Japan. Document SC/65a/SM12rev submitted to the 65a meeting of the Scientific Committee. Available online at: <http://iwc.int/sc65adocs>
33. Baulch, S. & Perry, C. 2013. Review of data available on the Dall's porpoise (*Phocoenoides dalli*) in Japanese waters. Document SC/65a/SM11 submitted to the 65a meeting of the Scientific Committee. Available online at: <http://iwc.int/sc65adocs>
34. Minamikawa, S., Shimada, H., Miyashita, T. and Moronuki, H. 2007. Estimates of Stock Levels for Six Cetacean Whaling Species from 1998-2001 Visual Inspection Data. Lecture Executive Summary Collection of the 2007 Japan Society of Fisheries Science Fall Conference. p151. In Japanese.
35. Ref 32, *ibid*.
36. IWC, 2001a. Annual Report of the International Whaling Commission, 53: 30-32
37. Ref 34, *ibid*. Fisheries Research Agency, 2012c. The State of International Fishing Resources in 2011 - Dall's porpoise.
38. Miyashita, T. 1991. Stocks and abundance of Dall's porpoises in the Okhotsk Sea and adjacent waters. Document submitted to the Scientific Committee, the 43rd IWC, Reykjavik, Iceland, May 1991. Document SC/43/SM7IWC, 2008 IWC, 2008. Annual report of the International Whaling Commission, 60: 31-33
39. Ref 9, *ibid*.
40. Perry, C. 2011. Tsunami stalls Japan's Dall's porpoise slaughter, but for how long? *The Ecologist*, November, 2011. Available online at: http://www.theecologist.org/blogs_and_comments/commentators/other_comments/1134726/tsunami_stalls_japans_dalls_porpoise_slaughter_but_for_how_long.html
41. EIA (2012) Scaled down Dall's porpoise hunt resumes in Japan. Available at: <http://www.eia-international.org/scaled-down-dalls-porpoise-hunt-resumes-in-japan>
42. Ref 40, *ibid*; Fujise, Y., Ishikawa, H., Saino, S. & Kawasaki, M. 1993. Catch and struck-and-lost rate in the Japanese Dall's porpoise hand-harpoon fishery. Report of the International Whaling Commission 42, 521-528. Kasuya, T. 2007. Japanese Whaling and other cetacean fisheries. *Environmental Science and Pollution Research* 10: 39-48.
43. Ref 40, *ibid*; Fujise, Y., Ishikawa, H., Saino, S. & Kawasaki, M. 1993. Catch and struck-and-lost rate in the Japanese Dall's porpoise hand-harpoon fishery. SC/44/SM18 in Report of the International Whaling Commission 43: 453-457.
44. Perry, 1999. Paper SC/51/SM46. Presented to the IWC Scientific Committee, May 1999. Fisheries Research Agency, 2012c. The State of International Fishing Resources in 2011 - Dall's porpoise.
45. Fisheries Research Agency, 2012c. The State of International Fishing Resources in 2011 - Dall's porpoise.
46. Ref 43, *ibid*. Wade, P. R., Bass, C. L. & Kasuya, T. 2008. A comparison of methods for calculating thresholds for evaluating levels of catch of Japan's Dall's porpoise (*Phocoenoides dalli*) hand-harpoon hunt. SC/60/SM24 presented to 60th annual meeting of the IWC Scientific Committee in Santiago, Chile.
47. Wade, P. R., Bass, C. L. & Kasuya, T. 2008. A comparison of methods for calculating thresholds for evaluating levels of catch of Japan's Dall's porpoise (*Phocoenoides dalli*) hand-harpoon hunt. SC/60/SM24 presented to 60th annual meeting of the IWC Scientific Committee in Santiago, Chile.
48. Ref 45, *ibid*.
49. Ref 45, *ibid*.
50. Okamura, H., Iwasaki, T. and Miyashita, T. 2008. Toward sustainable management of small cetacean fisheries around Japan. *Fisheries Science*, 74: 718-729
51. IWC, 1984. Annual Report of the International Whaling Commission, 34:57 IWC, 1991a. Annual Report of the International Whaling Commission, 41: 77-79 IWC, 1999a. Annual Report of the International Whaling Commission, 51: 36 IWC, 2001a. Annual Report of the International Whaling Commission, 53: 30-32 IWC, 2008. Annual report of the International Whaling Commission, 60: 31-33
52. Ref 13, *ibid*.
53. Ref 9, *ibid*. Funahashi, N. & Kasuya, T. 2012. Review of Baird's Beaked Whale (*Berardius bairdii*) around Japan, exploitation, current knowledge and suggestions for future research. SC/64/SM8 Presented to the IWC Scientific Committee, June 2012.
54. Ref 13, *ibid*.
55. IWC, 2012. Report of the scientific committee. Annex L: Report of the Sub-Committee on Small Cetaceans International Whaling. Available at <http://iwc.int/office/cache/downloads/arsnoxddh0080kcgsw4808/Annex-L-SM.pdf>
56. IWC, 1991. Annual Report of the International Whaling Commission, 41: 77-80
57. Ref 52, *ibid*.
58. Ref 13, *ibid*.
59. Taylor, B.L., Baird, R., Barlow, J., Dawson, S.M., Ford, J., Mead, J.G., Notarbartolo di Sciara, G., Wade, P. & Pitman, R.L. 2011. *Globicephala macrorhynchus*. In: IUCN 2012. IUCN Red List of Threatened Species. Version 2012.1. <www.iucnredlist.org>. Downloaded on 21 August 2012.
60. Ref 56, *ibid*.
61. IWC, 1987. Annual Report of the International Whaling Commission, 37: 51
62. Ref 12, *ibid*.
63. Ref 12, *ibid*.
64. Ref 12, *ibid*.
65. Ref 12, *ibid*.
66. Ref 12, *ibid*.
67. Ref 9 and 24, *ibid*; IWC, 1993a. Report of the scientific committee 43: 82-85
68. Ref 61, *ibid*.
69. Ref 2, *ibid*. Kasuya, T. 1985. Effect of exploitation on reproductive parameters of the spotted and striped dolphins off the Pacific coasts of Japan. *Scientific Report of the Whales Research Institute* 36, 107-138
70. Ref 61, *ibid*.
71. Ref 24, *ibid*.
72. Ref 24 and 9 *ibid*; Wells, R. S. and Scott, M. D. 1999. Bottlenose dolphin - *Tursiops truncatus* (Montagu, 1821) In: Handbook of marine mammals (Ridgway SH, Harrison SR, eds.) Vol. 6: The second book of dolphins and porpoises. pp. 137-182.
73. Ref 61, *ibid*.
74. Odell, D. K. and McClune, K. M. 1999. *Pseudorca crassidens*. In: Handbook of Marine Mammals (Ridgway SH, Harrison SR, eds.) Vol. 6: The second book of dolphins and porpoises. pp. 213-244.
75. Culik, B. 2010. Odontocetes. The toothed whales: "*Pseudorca crassidens*". UNEP/CMS Secretariat, Bonn, Germany. http://www.cms.int/reports/small_cetaceans/index.htm
76. Ref 24, *ibid*.
77. Elsa Nature Conservancy, 2007. Pacific white-sided dolphins. Available at <http://en.elsaenc.net/report/publiccomment/>.
78. Data is only complete for all species from 1979 onwards (Japan's progress reports to the IWC). Live captures are not included. Data on Baird's beaked whale catches are included from 1932 onwards (Funahashi & Kasuya, 2012) and for striped dolphin from 1942 onwards (Kasuya, 1999b) but these should not be considered to represent total catches of these species during these years due to incomplete reporting. Data from 1963-1978 are derived from Kishiro & Kasuya (1993) and Kasuya (1982). N/A indicates years when data are not available.
79. Ref 24, *ibid*. Funahashi, N. & Kasuya, T. 2012. Review of Baird's Beaked Whale (*Berardius bairdii*) around Japan, exploitation, current knowledge and suggestions for future research. SC/64/SM8 Presented to the IWC Scientific Committee, June 2012. Kasuya, T. 1999b. Examination of the reliability of catch statistics in the Japanese coastal sperm whale fishery. *Journal of Cetacean Research & Management* 1 (1) 109-121.
80. Kishiro, T. and Kasuya, T. 1993. Review of Japanese dolphin drive fisheries and their status. Report of the International Whaling Commission 43, 439-452. Kasuya, T. 1982. Preliminary report of the biology, catch and population of *Phocoenoides* in the western North Pacific, pp 3-19. In: Clarke JG (ed), *Mammals of the Sea*. FAO Fisheries Series, IV. FAO, Rome, 531 pp.
81. Ref 78, *ibid*. The PBR thresholds calculated by Funahashi & Baker (2011) use a recovery factor of 0.5. However they note that for some species it could be argued that a recovery factor of 0.1 is more appropriate, this would result in even lower thresholds.
82. Miyashita, T. 1991. Stocks and abundance of Dall's porpoise in the Okhotsk Sea and adjacent waters. SC/43/SM7. 24pp.
83. Miyashita, T., Iwasaki, T. and Moronuki, H. 2007a. An Estimate of Dall's/True's Porpoise Stock Levels in the Northwestern Pacific Ocean. Lecture Executive Summary Collection of the 2007 Japan Society of Fisheries Science Fall Conference. p.164. In Japanese.
84. PBR threshold calculated by Wade et al. (2008). Ref 45, *ibid*.
85. Ref 80, *ibid*.
86. Ref 81, *ibid*.
87. Ref 82, *ibid*.
88. Miyashita, T. 1986. Abundance of Baird's beaked whales off the Pacific coast of Japan. *Rep.Int.Whal. Comm.* 36: 383-386.
89. Miyashita, T. 1990. Population estimate of Baird's beaked whales off Japan. Paper IWC/SC/42/SM28, presented to the IWC Scientific Committee in 1990. (Available from IWC Secretariat).
90. Miyashita, T. and Kato, H. 1993. Population estimate of Baird's beaked whales off the Pacific coasts of Japan using sighting data collected by R/V Shunyu Maru, 1991 and 1992. Paper IWC/SC/45/SM6, presented to the IWC Scientific Committee in 1993. (Available from IWC Secretariat).
91. Okamura, H., S. Minamikawa, H.J. Skaug, and T. Kishiro. 2011. Abundance Estimation of Long-Diving Animals Using Line Transect Methods. *Biometrics*. DOI: 10.1111/j.1541-0420.2011.01689.x.
92. Funahashi, N. & Kasuya, T. 2012. Review of Baird's Beaked Whale (*Berardius bairdii*) around Japan, exploitation, current knowledge and suggestions for future research. SC/64/SM8 Presented to the IWC Scientific Committee, June 2012.
93. Ref 80, *ibid*.
94. Ref 87, *ibid*.
95. Ref 87, *ibid*.
96. Based on average catch for 2009 and 2010 only, as it is in these years that data are available on catch by sea area (Funahashi & Kasuya, 2012). Ref 89, *ibid*.
97. Calculated using the combined population estimated by Miyashita & Kato (1993) and Miyashita (1990). Ref 87 and 88 *ibid*.
98. IWC, 1992b. Report of the scientific committee. Annex G: Report of the Standing Sub-Committee on Small Cetaceans. 42: 178-221.
99. Ref 96, *ibid*.
100. Ref 12, *ibid*.
101. PBR threshold cited in Kanaji et al. 2011; Ref 12, *ibid*.
102. Miyashita, T. 1993. Abundance of dolphin stocks in the western North Pacific taken by the Japanese drive fishery. Reports of the International Whaling Commission 43: 417-437.
103. Ref 99, *ibid*.
104. Ref 99, *ibid*.
105. Ref 99, *ibid*.
106. Ref 99, *ibid*.
107. Ref 99, *ibid*.
108. Ref 99, *ibid*.
109. Ref 99, *ibid*.
110. Miyashita, T., Iwasaki, T. and Moronuki, H. 2007b. An Estimate of Pacific White-Sided Dolphin Stock Levels in the Vicinity of Japan from 1992-96 Visual Inspection Data. The Mammalogical Society of Japan 2007 Conference Program Collection of Executive Summaries. p.129. In Japanese.
111. Ref 3, *ibid*.
112. Wade, P. R., Reeves, R. R. and Mesnick, S. L. 2012. Social and Behavioural Factors in Cetacean Responses to Overexploitation: Are Odontocetes Less "Resilient" Than Mysticetes? *Journal of Marine Biology*, 195. doi:10.1155/2012/567276.
113. Fujise, Y., Ishikawa, H., Saino, S. & Kawasaki, M. 1993. Catch and struck-and-lost rate in the Japanese Dall's porpoise hand-harpoon fishery. SC/44/SM18 in Report of the International Whaling Commission 43: 453-457.
114. Ref 1, *ibid*.
115. Ref 14, *ibid*.
116. Maas, B. 2003. The potential stress effects of whaling operations and the welfare implications for hunted cetaceans. IWC/55/WK19. International Whaling Commission.
117. Ref 105, *ibid*.
118. Ref 105, *ibid*.
119. Aguilar, A., Borrell, A., Reijnders, P.J.H., 2002. Geographical and temporal variation in levels of organochlorine contaminants in marine mammals. *Marine Environmental Research* 53, 425-452.
120. ICES, 2010. Report of the Working Group on Marine Mammal Ecology (WGME), 12-15 April 2010, Horta, The Azores. ICES CM 2010/ACOM:24. 212 pp.
121. Ref 112, *ibid*. UNEP, 1998. Persistent organic pollutants. Information kit, Montreal.
122. Aguilar, A. and Borrell, A. 1994. Abnormally high polychlorinated biphenyl levels in striped dolphins (*Stenella coeruleoalba*) affected by the 1990-1992 Mediterranean epizootic. *Science of the Total Environment* 154: 237-24 Beland, P., De Guise, S., Girard, C., Lagace, A., Martineau, D., Michaud, R., Muir, D. C. G., Norstrom, R. J., Pelletier, E., Ray, S. and Shugart, L. R. 1993. Toxic compounds and health and reproductive effects in St. Lawrence beluga whales. *Journal of Great Lakes Research* 19, 766-775.
123. De Guise, S., Lagace, A., and Beland, P. 1994. Tumors in St Lawrence beluga whales (*Delphinapterus leucas*). *Veterinary Pathology* 31(4): 444-449. Hall, A.J., McConnell, B.J., Rowles, T.K., Aguilar, A., Borrell, A., Schwacke, L., Reijnders, P.J.H., and Wells, R.S. 2006. Individual-Based Model Framework to Assess Population Consequences of Polychlorinated Biphenyl Exposure in Bottlenose Dolphins. *Environmental Health Perspectives* 114(Suppl 1): 60-64 Jenson, P., Bennett, P.M., Deaville, R., Allichin, C.R., Baker, J.R., and Law, R.J. 2005. Relationships between polychlorinated biphenyls and health status in harbour porpoises (*Phocoena phocoena*) stranded in the UK. *Environmental Toxicology and Chemistry* 24(1): 238-248.
124. Kannan, K., Sinha, R. K., Tanabe, S., Ichihashi, H., Tatsukawa, R. 1993. Heavy metals and organochlorine residues in Ganges river dolphins from India. *Marine Pollution Bulletin* 26:159-162.
125. Lahvis, G. P., Well, R. S., Kuehl, D. W., Stewart, J. L., Rhinehart, H. L., & Via, C. S. 1995. Decreased lymphocyte responses in free-ranging bottlenose dolphins (*Tursiops truncatus*) are associated with increased concentrations of PCBs and DDT in

- peripheral blood. *Environmental Health Perspectives* 103: 62-72.
- Martineau et al., 1994 Martineau, D., De Guise, S., Fournier, M., Shugart, L., Girard, C., Lagace, A., Beland, P. 1994. Pathology and toxicology of beluga whales from the St. Lawrence Estuary, Quebec, Canada. Past, present and future. *Science of The Total Environment* 154(2-3): 201-215
- Reijnders & Brasseur, 1992; Reijnders, P. J. H. & Brasseur, S. M. 1992. Xenobiotic induced hormonal and associated developmental disorders in marine organisms and related effects in humans, an overview. In T. Colborn, & C. Clement (Eds.), *Advances in modern environmental toxicology* 21: 131-146.
- Reijnders, P.J.H., Aguilar, A. and Donovan, G.P. (eds). 1999. *Chemical Pollutants and Cetaceans. Journal of Cetacean Research and Management (Special Issue 1)*. Cambridge, UK.
- Simmonds, M. P. and Mayer, S. J. 1997. An evaluation of environmental and other factors in some recent marine mammal mortalities in Europe: implications for conservation and management. *Environmental Review* 5, 89-98
- Schwacke L.H., Voit E.O., Hansen L.J., Wells R.S., Mitchum G.B., Hohn A.A. et al. 2002. Probabilistic risk assessment of reproductive effects of polychlorinated biphenyls on bottlenose dolphins (*Tursiops truncatus*) from the southeast United States coast. *Environmental Toxicology and Chemistry* 21:2752-2764.
- Wells, R.S., Tornero, V., Borrell, A., Aguilar, A., Rowles, T.K., Rhinehart, H.L., Hofmann, S., Jarman, W.M., Hohn, A.A., and Sweeney, J.C. 2005. Integrating life-history and reproductive success data to examine potential relationships with organochlorine compounds for bottlenose dolphins (*Tursiops truncatus*) in Sarasota Bay, Florida. *Science of the Total Environment* 349(1-3): 106-119
117. De Guise et al. 1995; De Guise S., Martineau D., Béland P., Fournier M. 1995. Possible mechanisms of action of environmental contaminants on St. Lawrence beluga whales (*Delphinapterus leucas*). *Environmental Health Perspectives*, 103 (4): 73-77
- Hall, A.J., McConnell, B.J., Rowles, T.K., Aguilar, A., Borrell, A., Schwacke, L., Reijnders, P.J.H., and Wells, R.S. 2006. Individual-Based Model Framework to Assess Population Consequences of Polychlorinated Biphenyl Exposure in Bottlenose Dolphins. *Environmental Health Perspectives* 114(Suppl 1): 60-64
- Jepson, P.D., Tregezza, N. and Simmonds, M.P. 2008. Disappearing bottlenose dolphins (*Tursiops truncatus*) - is there a link to chemical pollution? *International Whaling Commission, SC/60/E*.
118. Weihe, P. & Joensen, H. D. 2012. Dietary recommendations regarding pilot whale meat and blubber in the Faroe Islands. *International Journal of Circumpolar Health*, Vol. 71.
119. Ref 8, *ibid.* JMHV, 1973. Provisional standard of mercury in fish and shell-fish, Director-General of Environmental Health Bureau. Japanese Ministry of Health and Welfare, Notification No. 99. JECFA, 2003. Joint FAO/WHO expert committee on food additives. 61st meeting. Rome. Kajiwara et al., 2002 Kajiwara, N., Watanabe, M., Tanabe, S., Nakamatsu, K., Amano, M., Myaziki, N. 2002. Specific accumulation and temporal trends of organochlorine contaminants in Dall's porpoises (*Phocoenoides dalli*) from Japanese coastal waters. *Marine Pollution Bulletin*, 44: 1089-1099
120. Endo, T., Haraguchi, K., Sakata, M., 2003. Renal toxicity in rats after oral administration of mercury-contaminated boiled whale livers marketed for human consumption. *Archives of Environmental Contamination and Toxicology* 44, 412-416
121. Ref 8, *ibid.*
122. Heldal, H. E., Foy, L. & Varskog, P. 2003. Bioaccumulation of 137 Cs in pelagic food webs in the Norwegian and Barents Sea. *Journal of Environmental Radioactivity*, 65(2): 177-185.
123. Ref 120, *ibid.*
124. Garnier-Laplace, J., Beaugelin-Seiller, K. & Hinton, T. G. 2011. Fukushima Wildlife Dose Reconstruction Signals Ecological Consequences. *Environ. Sci. Technol.* 45, 5077-5078.
125. <http://www.jfa.maff.go.jp/e/inspection/>
126. JMHV, 1973. Provisional standard of mercury in fish and shell-fish, Director-General of Environmental Health Bureau. Japanese Ministry of Health and Welfare, Notification No. 99.
127. Weihe, P. & Joensen, H. D. 2012. Dietary recommendations regarding pilot whale meat and blubber in the Faroe Islands. *International Journal of Circumpolar Health*, Vol. 71.
128. Ref 125, *ibid.*
129. Matsutani, M., 2010. Taiji locals test high for mercury: In surprise, experts fail to discover any signs of illness, *Japan Times*, May 10, 2010, p. 1. Matsutani, M. 2010. Most Taiji residents rest easy, refuse to change diet, *Japan Times*, May 10, 2010, p. 2.
130. Ref 3, *ibid.*
131. Moore, J.E., and Merrick, R., editors. 2011. *Guidelines for Assessing Marine Mammal Stocks: Report of the GAMMS III Workshop*, February 15 - 18, 2011, La Jolla, California. Dept. of Commerce, NOAA Technical Memorandum NMFS-OPR-47.
132. Ref 4, *ibid.*
133. Ministry of Agriculture, Forestry and Fisheries. 2002. *Substantial Progress for Japan - A Weekly update of news from the Japanese Ministry of Agriculture, Forestry and Fisheries*. Number 461 - July 12, 2002. Available at <http://www.maff.go.jp/mud/461.html> and <http://www.whales.org.au/govern/maff-54th.html>
134. Ref 2, 5, 6, 95 *ibid.* IWC, 1990a. Chairman's Report of the Forty-Second Meeting, Appendix 4. Resolution on the directed take of Dall's porpoises. Report of the International Whaling Commission, 41:48-9 IWC, 1990b. Annual Report of the International Whaling Commission, 40: 73-76 IWC, 1991b. Annual Report of the International Whaling Commission, 41: 80 IWC, 1993a. Report of the scientific committee, 43: 82-85 IWC, 1993b. Chairman's Report of the Forty-Fifth Annual Meeting, Appendix 10. Resolution on the directed take of striped dolphins. Report of the International Whaling Commission, 44:34. IWC, 1998a. Annual Report of the International Whaling Commission, 48: 106-112 IWC, 1999b. Chairman's Report of the Fifty-First Annual Meeting, Appendix 10. IWC Resolution 1999-9. Resolution on Dall's porpoise. Annual Report of the International Whaling Commission, 1999:55-6. IWC, 2001b. Chairman's Report of the Fifty-Third Annual Meeting. IWC Resolution 2001-12. Resolution on Dall's Porpoise.
135. Ref 3, *ibid.*





ENVIRONMENTAL INVESTIGATION AGENCY (EIA)

EIA - LONDON

62/63 Upper Street

London N1 0NY, UK

Tel: +44 (0) 20 7354 7960

Fax: +44 (0) 20 7354 7961

email: ukinfo@eia-international.org

www.eia-international.org



EIA - WASHINGTON, DC

PO Box 53343

Washington, DC 20009 USA

Tel: +1 202 483-6621

Fax: +1 202 986-8626

email: info@eia-global.org

www.eia-global.org